

Protocols for early Treatment COVID and potentially other “viral illness”.

I generally follow the McCullough protocol for COVID treatment. The optimal situation is to be treated within 5 days. For this reason, I keep both Hydroxychloroquine and Ivermectin in my house and you should too. You can read all the protocols at c19protocols.com. It is not one size fits all but depends on symptoms and your particular health etc. I would print off several protocols from that site to keep handy. You may need to take them to your physician.

I think, with time, we will discover that this whole spectrum from “COVID”, Vaccine effects, and Shedding damage is actually all the same issue-- “Spike Protein syndrome” in different forms, the initial contact (whether virus or contact poison) is the most deadly.

This is my protocol, and is not official medical advice for any individual. But if I were ill, this is what I would do for myself, and you can see the details below in the algorithm by Dr. McCullough.

If sick from “COVID”:

Make sure you are on the supplements outlined in tab under Preventive strategies. D, Zinc and others critical to success.

Begin hydroxychloroquine ASAP 400 mg BID (Twice a day) day 1, then 200 BID for 5-10 days. Azithromycin 250 mg BID first day 250 mg daily X 5-10 days. If you get too much nausea or diarrhea or are allergic to Azithromycin, Doxycycline 100 mg BID x 5-10 days.

In some cases Ivermectin .3-.4 mg/kg 5-7 days


Baby ASA (80 mg) day 1 till over disease.

If any pulmonary symptoms such as cough or Shortness of Breath (SOB), you may need an inhaler or nebulizer. SOB is a serious risk factor and needs competent medical attention. Some people have made a 50% Hydrogen Peroxide solution with a drop of iodine and used in nebulizer successfully. I use Budesonide nebulizer by prescription.

See next page for the entire McCullough algorithm .

For thoughts about shedding and vaccine remorse go to that tab.

Lee Merritt MD



BMI=body mass index, Dz=disease, DM=diabetes mellitus, CVD=cardiovascular disease, CKD=chronic kidney disease, yr=years, HCCQ=hydroxychloroquine, AZM=azithromycin, DOXY=doxycycline, IVM=ivermectin, VTE=venous thrombo-embolism, EUA=Emergency Use Authorization (U.S. administration)